

Multimorbidity and systematic reviews

Susan Smith

Department of General Practice, RCSI

HRB Centre for Primary Care Research

Inchicore Family Doctors





Conflict interest

- Primary research on interventions for people with multimorbidity
- Update Cochrane review due 18 months
- Seeking grant funding
- Current funding
 - Health Research Board Ireland
 - Health Services Executive, Ireland



Overview

- Multimorbidity (MM)
- Systematic review of interventions designed to improve outcomes for patients with MM
- Incorporating MM into existing systematic reviews
- MM and Guidelines



- 59 year old woman
- Living alone
- IHD; Depression; Neurological condition; **Arthritis**
- Multiple medications including warfarin

Presents with pain in her right shoulder





THE ELEPHANT IN THE ROOM





Multimorbidity vs Comorbidity






HRB Centre for Primary Care Research

Multimorbidity in primary care and general practice


Multimorbidity and age





Deprivation and multimorbidity

In Scotland, people living in more deprived areas develop multimorbidity 10 years before those living in the most affluent areas



Barnett et al. Lancet. 2012 Jul 7;380(9836):37-43



theguardian

News Sport Comment Culture Business Money Life & style 1

News > Society > NHS

NHS could be 'overwhelmed' by people with long-term medical conditions

One of health service's most senior figures warns there needs to be a serious rethink of how patients are cared for

Denis Campbell, health correspondent The Guardian, Friday 3 January 2014 19.30 GMT

Healthy patient £288 per year vs £2599 for person with three or more conditions (multimorbidity)



Key issues (BMJ editorial series 2012)

- "Ordering the chaos"
- Mental health
- Continuity
 - Relationship and information
- Managing medicines
- Need interventions and patient oriented outcomes
- Treatment burden



Multimorbidity and systematic reviews

- Reviews with MM as topic
- Incorporating MM into existing reviews



Interventions for improving outcomes in patients with multimorbidity in primary care and community settings: Systematic review

Susan M Smith¹, Hassan Soubhi², Martin Fortin², Catherine Hudon², Tom O'Dowd³ ¹HRB Centre for Primary Care Research, RCSI Medical School, Dublin ²Department of Family Medicine, University of Sherbrooke, Quebec ³Department of Public Health and Primary Care, Trinity College Dublin






HRB Centre for Primary Care Research

Cochrane systematic review within EPOC

- Studies
 - RCTs, CCTs, CBAs and ITS
- Participants
 - Two or more conditions
- Interventions
 - Any intervention designed to improve outcomes in individuals defined as having multimorbidity
 - Primary care and community settings
 - Classification: EPOC taxonomy
- Outcomes



Results: Search





Included studies

- Ten studies; all RCTs
 - 3407 patients
 - 8 in USA and 2 in UK
 - Majority 6-12 months
 - 8 included patients with broad range of conditions though elderly
 - 2 focused on co-morbidities
- Overall minimal risk of bias though consideration of contamination of control patients was generally inadequate



Results: Interventions

Interventions:

- 6 organisational
- 4 patient oriented

Multifaceted including:

- \circ Case management
- Enhanced skill mix in teams
- $_{\odot}$ Structured care provision
- Patient focussed approaches such as self-care and self-management



Intervention element	Study
1. Professional	
Health Educator	Eakin
Care manager (non-clinical)	Bognor
Clinical nurse managers	Boult, Katon, Lin, Sommers
Pharmacists	Krska
Social workers	Sommers
2. Financial	
No study	
3. Organisational	
Structured visits and/or care plans	Eakin, Bognor, Boult, Katon, Krska
Structured telephone contact	Eakin
Enhanced multidisciplinary team	Boult, Katon, Lin, Sommers
4. Patient oriented	
Self management support	Eakin, Boult, Lorig
Individual patient programme	Bognor, Boult
Patient education	Katon
Problem solving therapy	Lin
Peer support	Lorig

Results: overview

- Variation in participants and interventions
- Co-morbidity vs multimorbidity
 - Problems with definitions and overlap with frailty
 - May need different interventions for different groups
- Timescale
 - Improvements in medication related measures
- Targeting risk factors or specific functional difficulties may be more effective



Implications: Research

- Definition of multimorbidity challenging
- Searching and labelling:
 - MeSH term needed
- Outcomes
 - Generic across conditions
 - Physical functioning, quality of life, goal attainment
- Economic analyses needed



Systematic review conclusions

- Limited evidence with focus on co-morbid conditions or multimorbidity in older patients
- Need for clear definition of participants, and appropriate outcomes
- Suggestion that interventions may be more effective if targeted at specific risk factors of functional difficulties
- Need for well-designed intervention studies



Where does MM fit into existing reviews?

- Participants
- Interventions
- Outcomes
- Results



Participants

- Are they likely to have Co-M / MM ?

- Potential exclusions ?

 Balance between external validity and individualised patient-centred interventions*

*Fortin and Smith. Improving the external validity of trials for people with multiple chronic conditions. Journal Co-Morbidity. 2013, Vol 2



HRB Centre for Primary Care Research

Diabetes Co-morbidity

- Cohort of 424 patients with type 2 diabetes from RCT
- Results
 - 90% two or more conditions
 - 25% had five or more chronic conditions
 - 189 conditions
- Mismatch between self-report and chart review
- GP visits and medication numbers related to multimorbidity but not diabetes control



Information in Clinical Trials*

- 161 RCTs from 11 Cochrane Reviews on diabetes, heart failure, COPD and stroke
- 43.5% described the prevalence of any comorbidity among participants with the index disease
- Replicability of inclusion and exclusion criteria only fair
- Proportion exclusions for comorbidities: 0 55%
- Very uncommon assessment of whether comorbidities were potential modifiers of treatment effects



Interventions

- Possible different effect in MM patients?
 - Heterogeneity of treatment effect?

• Would intervention add to treatment burden in MM?



What outcomes matter in MM?

- Clinical Outcomes
 - Disease Specific Measures, Clinical quality measures, Risk factors, body weight, frailty/physical fitness
- Patient-reported Outcomes
 - Psychological, Behavior, Daily functioning, Social, Treatment burden, Shared Decision Making, Goal Setting, Satisfaction with care provision



- Health Care System
 - Health care utilization, processes of care, accessibility of services, safety
- Cost Outcomes
- Other Outcomes
 - Health care utilization, processes of care, accessibility of services, safety

Results

- ? Potential for sub-group analysis
 - Impact MM on intervention effect
 - How will you define MM
- Consideration of generalisability of review findings



Challenges

- Unclear reporting in original trials
- Search strategies
 - What if disease focused review?
- Study designs
 - Pragmatic trials with quasi-experimental designs may be more likely in MM



MM and Guidelines

• Mr B: 75 yr old with Diabetes and COPD

 Mrs A: A 78-year-old woman with previous MI, type 2 diabetes, osteoarthritis, COPD and depression

Hughes et al. Age and Ageing, 2013, 42(1)



MM and Guidelines

- Mr B: 75 yr old with Diabetes and COPD •
 - 5 medications (+8)
 - 6 self-care/lifestyle alterations
 - 5-6 routine primary care appointments
- Mrs A: A 78-year-old woman with previous MI, type 2 diabetes, ۲ osteoarthritis, COPD and depression
 - 11 medications (minimum, +10)
 - 9 self-care/lifestyle alterations
 - 8–10 routine primary care appointments + smoking cessation support and pulmonary rehabilitation if she chose to accept a referral.

Hughes et al. Age and Ageing, 2013, 42(1)



MM and Guidelines

- Increasing consensus on need to adapt guidelines to take account MM
- Balance between clinical utility and levels evidence
- Challenge is how to adapt them
 Cross referencing with electronic delivery



Summary

- MM common and important
- Is evidence to guide policy and future research
- Should be considered for all reviews
- Challenge for EBM, guidelines and clinical care delivery but need evidence to support decision making



Thank you

Acknowledgements: Prof Martin Fortin Prof Cynthia Boyd

